

Nguyen Van Quynh.

PhD in Physics and Mathematics, Lecturer

Faculty of Basic Science, Hanoi University of Industry, Ha Noi, Vietnam.

Le Anh Thang.

Master of Mathematical Sciences, Lecturer

Faculty of Basic Science, Hanoi University of Industry, Ha Noi, Vietnam.

THEOREM ON UNIFORM CONTINUITY OF CONONICAL POTENTIAL

Abstract: *Cononical potential $\int K_p(z; \zeta) d\gamma(\zeta)$ is important in the theory of subharmonic and δ -subharmonic functions. Classical properties were presented in many monographs, for example, in the works of N. S. Landkoff and V. S. Azarin. The paper considers the case: measure γ in the plane. For any $z \in \mathbb{C}$ we consider the Cononical potential as an element of the spaces $L_p(\gamma; \mathbb{C})$. In this article we give a sufficient condition on a measure γ the function $\int K_p(z; \zeta) d\gamma(\zeta) \in L_p(\gamma; \mathbb{C})$ to be uniformly continious in the parameter z in \mathbb{C} .*

Key words: *Cononical potential, Borel measure, uniform continuity, Minkowski inequality, Lebesgue measure.*

Нгуен Ван Куинь

Кандидат физико-математических наук, старший преподаватель
Факультета фундаментальной науки, университет Ханой
промышленности, Ha Noi, Вьетнам.

Ле Ань Тханг

Магистр математических наук, старший преподаватель
Факультета фундаментальной науки, университет Ханой
промышленности, Ha Noi, Вьетнам.

ТЕОРЕМА О РАВНОМЕРНОЙ НЕПРЕРЫВНОСТИ ЛОГАРИФМИЧЕСКОГО ПОТЕНЦИАЛА

Анотация: Кононический потенциал $\int K_p(z; \zeta) d\gamma(\zeta)$ играет важную роль в теории субгармонических и δ -субгармонических функций. Классические свойства были представлены во многих монографиях. Отметим, что А.Ф. Гришин и А. Шуиги изучили случай, когда мера γ есть ограничение меры Лебега на полуплоскости. В работе рассматривается случай: мера γ в плоскости. Для любого $z \in \mathbb{C}$ мы рассматриваем кононический потенциал как элемент пространств $L_p(\gamma; \mathbb{C})$. В этой статье приводится достаточное условие на меру γ для того, чтобы функция $\int K_p(z; \zeta) d\gamma(\zeta) \in L_p(\gamma; \mathbb{C})$ была равномерно непрерывной по параметру z в \mathbb{C} .

Ключевые слова: кононический потенциал, борелевская мера, равномерная непрерывность, неравенство Минковского, мера Лебега.

The theory of generalized functions is a basis for progress in many areas of mathematics and finds extensive applications in physics. One of the subdivisions of mathematics which is strongly influenced by the theory of generalized functions is the theory of subharmonic functions. Using generalized functions Azarin [1] constructed a theory of limit sets of subharmonic functions and measures, which is a significant contribution to the theory of subharmonic functions.

The best-known result in this area belong to Hormander [7]. In particular, he proved that the convergence of a sequence of subharmonic functions in a domain G as a sequence of the theory of generalized functions in G is equivalent to the convergence in the space $L_{m,loc}(\gamma; \mathbb{C})$.

We will use the following notation:

$$B(z_0; R) = \{z \in \mathbb{C} : |z - z_0| \leq R\}; CB(z_0; R) = \mathbb{C} / B(z_0; R).$$

In the theory of subharmonic and δ -subharmonic functions in the plane \mathbb{C} , an important role is played by the kernel:

$$K_p(z; \zeta) = \operatorname{Re} \left(\ln \left(1 - \frac{z}{\zeta} \right) + \frac{z}{\zeta} + \dots + \frac{1}{p} \frac{z^p}{\zeta^p} \right),$$

where $p \in \mathbb{N}$. For all $z, \zeta \in \mathbb{C}$ we have the inequality ([8], Lemma 2):

$$|K_p(z; \zeta)| \leq M(p) \frac{|z|^p}{|\zeta|^p} \min \left\{ 1; \frac{|z|}{|\zeta|} \right\}, \quad (1)$$

where $M(p)$ depends only on p . Let γ be the Radon measure in \mathbb{C} . We consider the following potential

$$K(z) = \int_C K_p(z; \zeta) d\gamma(\zeta),$$

which we shall call the canonical potential of the measure γ .

In this paper, we consider this Canonical potential $K(z)$ as a map from the space \mathbb{C} into the space $L_p(C; d\gamma(\zeta))$. In this case, we can write $K(z) : C \rightarrow L_p(\gamma)$.

The convergence of a sequence $u_n(z)$ to $u(z)$ in the space $L_m(\gamma; C)$ means the following. If γ is the restriction of the Lebesgue measure to a compactum K contained in the space \mathbb{C} , then $\iint |u_n(z) - u(z)| d\gamma(z) \rightarrow 0 \quad (n \rightarrow \infty)$.

We state several well-known results obtained in this area. We have already mentioned Hormander's proposition [8]. Hormander considered the multidimensional case; we consider the case where the dimension of the space is $m=2$, although the requirement $m=2$ is not essential in many of our arguments. It is important for what follows that the dependence of $K(z)$ on z be uniformly continuous.

In the present paper we find sufficient conditions on a measure γ with support compactly embedded in G that guarantee the convergence in the space $L_m(\gamma; C)$ of sequences of subharmonic functions that converge in the sense of the theory of generalized functions.

We state several well-known results obtained in this area. We have already mentioned Hormander's proposition. Hormander considered the multidimensional case; we consider the case where the dimension of the space is $m=2$, although the requirement $m=2$ is not essential in many of our arguments. The following was proved in [10], Theorem 5. Suppose that γ_n is a uniformly compactly supported sequence of Borel measures, $\text{supp } \gamma_n \subset \mathbb{C}$, and the measures γ_n have uniformly bounded variation and weakly converge to the zero measure.

The uniform continuity of these functions is a restriction on the measure γ which appears in the theorems that are the principal results of the paper. It is important for what follows that the dependence of $K(z)$ be uniformly continuous.

Theorem. Let $m > 1$ be an arbitrary fixed number. Suppose that γ is a positive finite Borel measure with compact support that satisfies the condition $\text{supp } \gamma \subset \mathbb{C}$. In addition, suppose that

$$\sup \left\{ \int_{B(z;\delta)} |K_p(z;\zeta)|^m d\gamma(\zeta) : z \in C \right\} \rightarrow 0; \quad (2)$$

$(\delta \rightarrow 0).$

Then the function $K(z): C \rightarrow L_m(\gamma)$ is uniformly continuous with respect to the variable z in the space \mathbb{C} .

Proof. We divide the proof into several stages.

1. Since we have

$$K_p(z;\zeta) = \text{Re} \left(\ln \left(1 - \frac{z}{\zeta} \right) \right) + \text{Re} \left(\frac{z}{\zeta} + \dots + \frac{1}{p} \frac{z^p}{\zeta^p} \right) = \ln \left| 1 - \frac{z}{\zeta} \right| + K_p^{-1}(z;\zeta)$$

Obviously the function $K_p^{-1}(z;\zeta): C \rightarrow L_m(\gamma)$ is uniformly continuous with respect to the variable z in the space \mathbb{C} .

2. We set

$$\psi(z_1; z_2) = \left(\int_C |\ln |z_1 - \zeta| - \ln |z_2 - \zeta||^p d\gamma(\zeta) \right)^{1/p}; \quad F(z) = \left(\int_C |\ln |z - \zeta||^p d\gamma(\zeta) \right)^{1/p}.$$

Since the Logarithmic potential as a function from \mathbb{C} into \mathbb{R} , is not bounded, we need to prove the convergence of the integrals introduced above.

We do this at the first stage. It follows from the Minkowski inequality that

$$F(z) \leq \left(\int_{B(z; \delta)} |\ln |z - \zeta||^p d\gamma(\zeta) \right)^{1/p} + \left(\int_{CB(z; \delta)} |\ln |z - \zeta||^p d\gamma(\zeta) \right)^{1/p} = J_1 + J_2.$$

where $CB(z; \delta) = C \setminus B(z; \delta)$.

We prove that the inequality

$$|\ln |z - \zeta|| \leq \ln \frac{1}{\delta} \quad (3)$$

holds for sufficiently small δ and for $\zeta \in (\text{supp } \gamma) \cap CB(z; \delta)$. From the inequality

(3) we obtain that $J_2 \leq \ln \frac{1}{\delta} (\gamma(C))^{1/p}$. By the hypothesis of the theorem, $J_1 \leq 1$ for

sufficiently small δ . Hence, $F(z)$ is finite. What has been proved can also be stated as follows: for any $z \in \mathbb{C}$ the function $K(z)$ is an element of the space $L_p(\gamma)$. The inequality $\psi(z_1; z_2) \leq F(z_1) + F(z_2)$ implies that $\psi(z_1; z_2)$ is finite.

3. We prove that the inequality

$$|\nabla_\zeta \ln |z - \zeta|| \leq \frac{1}{\delta}. \quad (4)$$

holds for $|z - \zeta| \geq \delta$.

$$\text{Let } z = x + iy; \zeta = \eta + i\tau \text{ we have } (\ln |z - \zeta|)'_\eta = -\frac{x - \eta}{|z - \zeta|^2}; (\ln |z - \zeta|)'_\tau = -\frac{y - \tau}{|z - \zeta|^2}.$$

From these estimates it is easy to obtain inequality

$$|\nabla_\zeta \ln |z - \zeta|| = \left((\ln |z - \zeta|)'_\eta \right)^2 + \left((\ln |z - \zeta|)'_\tau \right)^2 = \frac{1}{|z - \zeta|} \leq \frac{1}{\delta}.$$

Thus the inequality (4) is proved.

4. This is the most essential part of the proof. We claim that

$$\psi(z; z_0) \rightarrow 0 \quad (z \rightarrow z_0).$$

Let δ be an arbitrary number. We assume that the inequality $|z - z_0| \leq \delta$ holds. Consecutive application of the Minkowski inequality [4] and the inclusion $B(z_0; 2\delta) \subset B(z; 3\delta)$ yields

$$\begin{aligned} \psi(z; z_0) \leq & \left(\int_{CB(z; 3\delta)} |\ln |z - \zeta||^p d\gamma(\zeta) \right)^{1/p} + \left(\int_{CB(z_0; 2\delta)} |\ln |z - \zeta| - \ln |z_0 - \zeta||^p d\gamma(\zeta) \right)^{1/p} + \\ & + \left(\int_{CB(z_0; 2\delta)} |\ln |z_0 - \zeta||^p d\gamma(\zeta) \right)^{1/p} = J_1 + J_2 + J_3. \end{aligned}$$

If $\zeta \notin B(z_0; 2\delta)$ and $|z - z_0| \leq \delta$, then the inequalities $|w - \zeta| > \delta$ and

$$|\nabla_\zeta \ln |z - \zeta|| \leq \frac{1}{\delta} \quad \text{hold for any } w \in B(z_0; |z - z_0|).$$

The estimate of the gradient implies that

$$|\ln |z - \zeta| - \ln |z_0 - \zeta|| \leq \frac{1}{\delta} |z - z_0|; J_2 \leq \frac{1}{\delta} (\gamma(C))^{1/p} |z - z_0|.$$

From the hypothesis of the theorem we obtain that

$$J_1 \rightarrow 0; \quad J_3 \rightarrow 0 \quad (\delta \rightarrow 0),$$

which proves the required assertion.

The collection of assertions that we have obtained is contradictory. Thus the theorem is proved.

Example. Let γ be a Lebesgue measure in the space \mathbb{C} . We will use the standard notation $d\gamma = dx dy$. Consider the integral in the condition (2) and, making a parallel translation, we obtain the equality for $\zeta = x + iy$:

$$J_1 = \int_{B(z; \delta)} |\ln |z - \zeta||^p dx dy = \int_{B(0; \delta)} |\ln |\zeta||^p dx dy.$$

$$\text{We introduce polar coordinates: } \begin{cases} x = r \cos \varphi \\ y = r \sin \varphi \end{cases}.$$

$$\text{We obtain the equality: } J_1 = \int_0^{2\pi} \int_0^\delta r |\ln r|^p dr d\varphi = 2\pi \int_0^\delta r |\ln r|^p dr.$$

It is obvious that condition holds for arbitrary p : $J_1 \rightarrow 0$; $(\delta \rightarrow 0)$.

References

1. N.S. Landkof (1966), Foundations of modern potential theory, GRFML, Science.
2. Azarin V. S. (2009), Growth theory of subharmonic functions Birkhanser, Basel, Boston, Berlin.
3. A.F. Grishin, Nguyen Van Quynh, I.V. Poedintseva (2014), Representation theorems of δ -subharmonic Functions, Visnyk of V.N.Karazin Kharkiv National University. Ser. "Mathematics, Applied Mathematics and Mechanics", No.1133. – P. 56-75.
4. Kadets, V.M. (2006), A course of Functional Analysis, Kharkov National University.
5. J Van Quynh Nguyen (2015), Various Types of Convergence of Sequences of Subharmonic Functions, Zh. Mat. Fiz. Anal. Geom, Volume 11, Number 1, 63–74.
6. A. F. Grishin, A. Chouigui (2008), Various types of convergence of sequences of δ -subharmonic functions, Math. sbornik. No. 199, 27–48.
7. L. Hörmander (1983), The Analysis of Linear Partial Differential Operators. II. Differential Operators with Constant Coefficients. Springer-Verlag, Berlin, Heidelberg, New York, Tokyo, 1983.
8. Levin B.Ya. Distribution of zeros of entire functions. / B.Ya. Levin – M.: GITTL. - 1956. - 632 p.
9. Nguyen Van Quynh, Theorem on uniform continuity of Logarithmic potential, Visnyk of science and education, Issue 5 (59), 6-10p.